

**OXFORD DICTIONARY OF**  
**BIOCHEMISTRY AND**  
**MOLECULAR BIOLOGY**  
**REVISED EDITION**

Managing Editor **Dr A D Smith** University College London

General Editors **Professor S P Datta** University College London  
**Dr G H Smith** University College London  
**Professor P N Campbell** (Chairman) University College London  
**Dr R Bentley** University of Pittsburgh  
**Dr H A McKenzie** Australian Defence Force Ac.

Subject Editors **Dr D A Bender** University College London  
**Dr A J Harris** University of Queensland  
**Professor T W Goodwin** University of Liverpool  
**Dr H A McKenzie** Australian Defence Force Ac.  
**Dr J H Parish** University of Leeds  
**Dr C Stanford** University College London

**EXHIBIT A**

**OXFORD**  
UNIVERSITY PRESS

# Editors

## General Editors

Dr A. D. Smith (Managing Editor) *Emeritus Reader in Biochemistry, and Honorary Research Fellow, Department of Molecular Pathology, University College London Medical School*

Professor S. P. Datta *Emeritus Professor of Medical Biochemistry, and Honorary Research Fellow, Department of Biochemistry and Molecular Biology, University College London*

Dr G. Howard Smith *Honorary Research Fellow, Department of Biochemistry and Molecular Biology, University College London*

Professor P. N. Campbell *Emeritus Professor of Biochemistry, and Honorary Research Fellow, Department of Biochemistry and Molecular Biology, University College London*

Professor R. Bentley *Professor Emeritus of Biochemistry, Department of Biological Sciences, University of Pittsburgh*

Dr H. A. McKenzie *Fellow, School of Chemistry, University of New South Wales, Australian Defence Force Academy, and Visitor, Molecular Medicine, John Curtin School of Medical Research, The Australian National University, Canberra*

## Subject Editors

Dr D. A. Bender *Senior Lecturer, Department of Biochemistry and Molecular Biology, University College London*

Dr A. J. Carozzi *NH and MRC Research Officer, Centre for Molecular and Cellular Biology, University of Queensland*

Professor T. W. Goodwin FRS *Emeritus Johnson Professor of Biochemistry, University of Liverpool*

Dr J. H. Parish *Senior Lecturer, School of Biochemistry and Molecular Biology, University of Leeds*

Dr S. C. Stanford *Senior Lecturer, Department of Pharmacology, University College London*

OXFORD

UNIVERSITY PRESS

Great Clarendon Street, Oxford OX2 6DP

Oxford University Press is a department of the University of Oxford.  
It furthers the University's objective of excellence in research, scholarship,  
and education by publishing worldwide in

Oxford New York

Athens Auckland Bangkok Bogotá Buenos Aires Calcutta  
Cape Town Chennai Dar es Salaam Delhi Florence Hong Kong Istanbul  
Karachi Kuala Lumpur Madrid Melbourne Mexico City Mumbai  
Nairobi Paris São Paulo Singapore Taipei Tokyo Toronto Warsaw

with associated companies in Berlin Ibadan

Oxford is a registered trade mark of Oxford University Press  
in the UK and in certain other countries

Published in the United States  
by Oxford University Press Inc., New York

© The General Editors, 1997

The authors have asserted their right under the Copyright, Designs and Patents Act, 1988,  
to be identified as authors of this work

First published 1997

Revised edition 2000

All rights reserved. No part of this publication may be reproduced,  
stored in a retrieval system, or transmitted, in any form or by any means,  
without the prior permission in writing of Oxford University Press,  
or as expressly permitted by law, or under terms agreed with the appropriate  
reprographics rights organization. Enquiries concerning reproduction  
outside the scope of the above should be sent to the Rights Department,  
Oxford University Press, at the address above

You must not circulate this book in any other binding or cover  
and you must impose this same condition on any acquirer

A catalogue record for this book is available from the British Library

Library of Congress Cataloging in Publication Data

(Data applied for)

ISBN 0 19 850673 2

Typeset by Market House Books Ltd, Aylesbury

Printed in Great Britain by  
Butler & Tanner Ltd, Frome

No  
G  
ini  
an  
we  
Di  
th  
ch  
jec  
pr  
me  
  
an  
na  
ab  
ex  
of  
wa  
sp  
tio  
bo  
co  
an  
ch  
or  
ha  
ter  
so  
to  
ary  
ref  
no  
spr  
tio  
var  
bec  
nu  
  
the

**enantiomorphic**

are mirror images of each other and thus are not superposable. 2 either of the two crystalline forms exhibited by a pair of enantiomers. Use of the term to mean **enantiomer** is deprecated.

**enantiomorphic** or **enantiomorphous** of, or pertaining to, an **enantiomorph**; pertaining to the phenomenon of, or displaying **enantiomorphism**. The term is often used synonymously with **enantiomeric** (enantiomeric molecules frequently form enantiomorphic crystals).

**enantiomorphism** the phenomenon of being related as between an object and its nonsuperposable mirror image. The term is used especially in relation to enantiomorphic crystals.

**enantiomorphous** see **enantiomorphic**.

**enantiotopic** 1 when chemically-like ligands in constitutionally equivalent locations (generally the two *a* ligands in *Caabc*) are related by a centre or plane of symmetry, or by an alternating axis of symmetry (but not by a simple axis of symmetry), they are enantiotopic. The two ligands are in a stereochemically different, mirror-image environment. If each *a* ligand of *Caabc* is replaced separately by a different, achiral ligand, *d*, the products are the two enantiomers of *Cabcd*. Example: the methylene hydrogens of ethanol are enantiotopic; if ethanol is written as a Fischer projection structure with OH at the top, H-C-H in the middle, and CH<sub>3</sub> at the bottom, the left-hand hydrogen of the central methylene is H<sub>S</sub>, while that at the right is H<sub>R</sub> (see *pro-R/pro-S convention*). Replacement of <sup>1</sup>H<sub>R</sub> by <sup>2</sup>H yields (+)-(R)-[1-<sup>2</sup>H]ethanol and the same replacement of <sup>1</sup>H<sub>S</sub> yields the enantiomer, (-)-(S)-[1-<sup>2</sup>H]ethanol. In another important compound, citric acid, the two CH<sub>2</sub>COOH groups are also enantiotopic. 2 the two faces of a double bond or of a planar cyclic ring system that are related by a symmetry plane but not by a C<sub>2</sub> axis (i.e., a two-fold axis of symmetry) are enantiotopic; the two faces show stereochemically different, mirror-image related environments. Separate addition of the same achiral reagent to the two faces (see *Re/Si convention*) gives enantiomeric products. Example: the simple addition of HCN to CH<sub>3</sub>-CHO yields a racemic mixture of the (R) and (S) cyanohydrins, CH<sub>3</sub>-CH(OH)-CN, with both faces of C=O being involved. The reduction of the C=O bond of CH<sub>3</sub>-CHO to form ethanol by alcohol dehydrogenase requires addition of a hydride ion from NADH at the C atom and a hydron at the O atom. Thus, reduction of CH<sub>3</sub>-CHO with NAD<sup>2</sup>H at its *A* face (see *diastereotopic* (def. 2)) yields (R)-[1-<sup>2</sup>H]ethanol and reduction of CH<sub>3</sub>-C<sup>2</sup>HO with *A*-NADH yields (S)-[1-<sup>2</sup>H]ethanol. The enzymatic reduction is stereospecific and only one of the enantiotopic faces of C=O is attacked; it is the same one (the *Re* face) in both of these situations. Compare *diastereotopic*.

**encapsidate** to surround (a particle of viral nucleic acid) with a **capsid**. —**encapsidation** *n*.

**encapsis** the association of myofibrils into bundles and the further association of these bundles into larger bundles, etc.

**encephalin** a variant spelling of **enkephalin**.

**encephalitis** inflammation of the brain.

**3' end** the end of a linear polynucleotide strand at which the 3'-hydroxyl group of the terminal nucleoside residue is normally not phosphorylated.

**5' end** the end of a linear polynucleotide strand at which the 5'-hydroxyl group of the terminal nucleoside residue is normally phosphorylated.

**endo+** a variant form of **endo+** (sometimes before a vowel).

**end capping** (in chromatography) the blocking of residual silanol groups on the surface of silica where these remain exposed after the bonding of C<sub>18</sub> or other alkyl chains to the silica in the formation of reversed-phase stationary phases for column chromatography. For this purpose hydrocarbyl silanes (see **silane** (def. 3)) having small alkyl (usually methyl) groups are used so that they can penetrate between the main bonded-phase groups.

**endemic** present in or peculiar to a more or less localized area, e.g. an endemic disease. Compare **enzootic**.

**endergonic** describing a process or reaction on which work must be done, i.e. one requiring an energy input, for it to take place. At constant pressure and temperature the free energy content of such a system increases. Compare **exergonic**. [From **endo+** plus Greek *ergon*, work.]

**end group** any residue at an extremity of a branched or linear macromolecule.

**end-group analysis** determination of both the nature and the number of terminal groups in a macromolecule, e.g. in proteins, the N- and C-terminal amino-acid residues; in polynucleotides, the 3'- and 5'-terminal nucleotide residues.

**endo+** or (sometimes before a vowel) **end+** comb. form meaning within, inner, absorbing, containing. Compare **exo+**. See also **intra+**.

**endo-** prefix (in chemical nomenclature) denoting insertion (of the additional constituent(s) specified) into the structure of (a named compound); e.g. endo-4a-glycine-[5-leucine]enkephalin; endo-Gly<sup>1a</sup>-[Leu<sup>2</sup>]enkephalin: Tyr-Gly-Gly-Phe-Gly-Leu: a synthetic polypeptide in which a glycine residue has been inserted between residues 4 and 5 of [5-leucine]enkephalin.

**endo-** prefix (in stereochemistry). See **conformation**.

**endoamylase** any **amylase** that hydrolyses nonterminal glycosidic linkages; it is a subcategory of **endoglycosidase**.

**endocrine** 1 describing or relating to any gland or other group of cells that synthesizes **hormones** and secretes them directly into the blood, lymph, or other intercellular fluid. 2 describing or relating to a secretion of endocrine tissue. 3 a secretory product of endocrine tissue; a hormone. Originally known as **internal secretion**. Compare **exocrine**.

**endocrine gland** or **ductless gland** any of the ductless glandular structures that secrete (one or more) hormones directly into the bloodstream.

**endocrinology** the science concerned with the endocrine organs, their products, and the effects of these products. — **endocrinological** *adj*.

**endocytic** 1 situated within a living cell but not belonging to the cell itself; intracellular. 2 an alternative term for **endocytotic** (see **endocytosis**).

**endocytosis** the uptake of external materials by cells through the mechanism of **phagocytosis** or **pinocytosis**. The term is often used interchangeably with **pinocytosis**. Compare **exocytosis**, **transcytosis**. See also **internalize**, **viropexis**. — **endocytic** or **endocytotic** *adj*.; — **endocytose** *vb*.

**endocytotic vesicle** see **pinocytotic vesicle**.

**endodeoxyribonuclease** see **deoxyribonuclease**.

**endoenzyme** 1 any intracellular enzyme. Compare **ectoenzyme**, **exoenzyme** (def. 1). 2 any enzyme that catalyses **endohydrolysis**. It may be an **endoglycosidase**, an **endonuclease**, or an **endopeptidase**. Compare **exoenzyme** (def. 2).

**endogenous** arising or developing within an organism, tissue, or cell, and excluding any consequences of externally added agents or materials. — **endogenously** *adv*.

**endoglin** a major glycoprotein of vascular endothelium that may be important in the binding of endothelial cells to integrins. It forms a heteromeric complex with the signalling receptors for transforming growth factor β (TGF-β). It has an RGD integrin-recognition motif and is a homodimer of disulfide-linked subunits. Example (precursor) from *Sus scrofa*: data-base code EGLN\_PIG, 653 amino acids (70.20 kDa).

**endoglycosidase** any enzyme within subclass EC 3.2, glycosidases, that hydrolyses nonterminal glycosidic linkages in oligo- or polysaccharides. Many activities of this type are known, e.g. from *Flavobacterium meningosepticum*.

**endohormone** any hormone acting within the individual organism that produces it. Compare **ectohormone**.

**endohydrolysis** the hydrolysis, esp. by an **endoenzyme**, of any linkage between residues in a biopolymer. For example, **endopeptidases** attack neither the C-terminal nor the N-terminal peptide linkages of an oligo- or polypeptide, and **endoglycosidases** attack the terminal glycosidic linkages at either the reducing or nonreducing end of an oligo- or polysaccharide.